

McLaren Print System Order

Order No: 41789
Order Date: 2019-01-09
User: Judy Fago
Phone: 586-493-3610

Ship Location: Gratiot Medical Bldg, Multi Specialty
36500 Gratiot, Suite 202
Clinton Twp, MI 48035

Forms

Quantity: 1000
Paragon Dept No: 60320
Dept Name: Family First
Company Number: 260

Order Total Price: 67.00

Item Number: MO-3359
Item Description: Normal Lab Results Card (Adult)
Revision Date: 1/2017
Print: 2 sided black and white
Paper: 65# White Cover
Size: 8.5 x 11
Fold: Bi-Fold (1/2)
Finish:
Drill:
Misc Info:

- Total cholesterol should be under 200
- HDL (good cholesterol) should be over 40 and can be improved with regular exercise.
- LDL (bad cholesterol) should be less than 130 for those at average risk. For people with moderate risk it should be less than 160. For people with coronary artery disease, diabetes, stroke, or cerebrovascular disease, LDL should be under 100. To lower it, eat foods low in fat and cholesterol in your diet and exercise regularly.
- Triglycerides (a blood fat) should be less than 150. Reduction of dietary carbohydrates (bread, pasta, cereal, fruit) and reduction of alcohol reduce triglycerides.
- Hemoglobin (Hgb) is a measure of red blood cells. If low, you have anemia.
- Normal fasting blood sugar or glucose is 70-100. Glucose tolerance if from 100-125, it may indicate diabetes.
- High H C (CMB) is a measure reflecting the average blood sugar over the last three months.
- CMB under 7.0 is recommended for diabetic patients.
- TSH (Thyroid screening test) is elevated in underactive thyroid and low in overactive thyroid.
- PSA (Prostate specific antigen) is a measure of prostate activity. Normal values are 0-4.0. Higher levels may require further testing.
- BUN and Creatine are kidney function tests when either are high, the kidneys are not functioning optimally.
- Electrolyte abnormalities may be caused by water pills, kidney problems, dietary deficiencies or other medical problems.
- Prostate Specific Antigen is a test for hidden blood in the stool, and one way to screen for colon cancer. If positive, you may need a colonoscopy.
- White Blood Cell Count is often elevated in patients with infections.
- PSA Routine pap smears are an important way to screen for cervical cancer. For women between the ages of 21 and 65, the American Cancer Society recommends a pap smear every one to three years.
- Mammogram: Please remember that about 40% of cancers cannot be found by mammograms. Early detection requires a combination of self-examination, yearly clinical breast exams, and yearly mammograms.

Thank you
for choosing

MACOMB
and the providers and staff
who serve you.

Enclosed are the results of
your recent laboratory tests.
for your personal file.

Please bring all medications to
your next provider appointment.

Spec Info: