

McLaren Print System Order

Order No: 47826 Reprint Previous Order No: 17647
 Order Date: 2019-08-19
 User: Kim Brass
 Phone: 8103001855

Ship Location: MMG Marysville Internal Med
 1750 Busha Hwy
 Marysville, MI 48040

Forms

Quantity: 100
 Paragon Dept No: 66300
 Dept Name: 66300
 Company Number: 810

Order Total Price: 0.00

Item Number: MM-320
 Item Description: Hib
 Revision Date: 04/2015
 Print: 2 sided black and white
 Paper: 20# White Text
 Size: 8.5 x 11
 Fold:
 Finish: None
 Drill: None
 Misc Info: ds; black and white

VACCINE INFORMATION STATEMENT

Hib Vaccine

What You Need to Know

(Haemophilus influenzae Type b)

Many Vaccine Information Statements are available in Spanish and other languages. For more information, visit www.hhs.gov/opa.

1 Why get vaccinated?

Haemophilus influenzae type b (Hib) disease is a serious disease caused by bacteria. It usually affects children under 5 years old. It can also affect adults with certain medical conditions.

Your child can get Hib disease by being around other children or adults who may have the bacteria and not know it. The germs spread from person to person. If the germs stay in the child's nose and throat, the child probably will not get sick. But sometimes the germs spread into the lungs or the bloodstream, and then Hib can cause serious problems. This is called invasive Hib disease.

Before Hib vaccine, Hib disease was the leading cause of bacterial meningitis among children under 5 years old in the United States. Meningitis is an infection of the lining of the brain and spinal cord. It can lead to brain damage and death. Hib disease can also cause:

- pneumonia
- severe swelling in the throat, making it hard to breathe
- infections of the blood, joints, bones, and covering of the heart
- death

Before Hib vaccine, about 20,000 children in the United States under 5 years old got Hib disease each year, and about 7% - 9% of them died.

Hib vaccine can prevent Hib disease. Since use of Hib vaccine began, the number of cases of invasive Hib disease has decreased by more than 99%. Many more children would get Hib disease if we stopped vaccinating.

2 Hib vaccine

Several different brands of Hib vaccine are available. Your child will receive either 3 or 4 doses, depending on which vaccine is used.

Doses of Hib vaccine are usually recommended at these ages:

- First Dose: 2 months of age
- Second Dose: 4 months of age
- Third Dose: 6 months of age (if needed, depending on brand of vaccine)
- Final Booster Dose: 12-15 months of age

Hib vaccine may be given at the same time as other vaccines.

Hib vaccine may be given as part of a combination vaccine. Combination vaccines are made when two or more types of vaccine are combined together into a single shot, so that one vaccination can protect against more than one disease.

Children over 5 years old and adults usually do not need Hib vaccine. But it may be recommended for older children or adults with spleen or sickle cell disease, before surgery to remove the spleen, or following a bone marrow transplant. It may also be recommended for people 1 to 18 years old with HIV. Ask your doctor for details.

Your doctor or the person giving you the vaccine can give you more information.

U.S. Department of Health and Human Services
 Centers for Disease Control and Prevention