

McLaren Print System Order

Order No: 49553 Reprint Previous Order No: 17840
Order Date: 2019-10-17
User: Alicia Mullett
Phone: 9893932850

Ship Location: MCLAREN OCCUPATIONAL HEALTH
801 Joe Mann Blvd Suite A 1st Floor
Midland , MI 48642

Forms

Quantity: 100
Paragon Dept No: 56052
Dept Name: occupational health
Company Number: 810

Order Total Price: 0.00

Item Number: MM-334
Item Description: Tdap (Tetanus, Diptheria, Pertussis)
Revision Date: 2/2015
Print: 2 sided black and white
Paper: 20# White Text
Size: 8.5 x 11
Fold:
Finish: None
Drill: None
Misc Info: ds; black and white

VACCINE INFORMATION STATEMENT

Tdap Vaccine

What You Need to Know

(Tetanus, Diphtheria and Pertussis)

More Vaccine Information Statements are available in Spanish and other languages. For more information, visit www.cdc.gov/vaccines/imz/downloads/011115-0001.pdf.

© 2015 U.S. Department of Health and Human Services. All rights reserved. For more information, visit www.cdc.gov/vaccines/imz/downloads/011115-0001.pdf.

1 Why get vaccinated?

Tetanus, diphtheria and pertussis are very serious diseases. Tdap vaccine can protect us from these diseases. And, Tdap vaccine given to pregnant women can protect newborn babies against pertussis.

TETANUS (Lockjaw) is rare in the United States today. It causes painful muscle tightening and stiffness, usually all over the body.

- It can lead to tightening of muscles in the head and neck so you can't open your mouth, swallow, or sometimes even breathe. Tetanus kills about 1 out of 10 people who are infected even after receiving the best medical care.

DIPHTHERIA is also rare in the United States today. It can cause a thick coating to form in the back of the throat.

- It can lead to breathing problems, heart failure, paralysis, and death.

PERTUSSIS (Whooping Cough) causes severe coughing spells, which can cause difficulty breathing, vomiting and disturbed sleep.

- It can also lead to weight loss, incontinence, and rib fractures. Up to 7 in 100 adolescents and 9 in 100 adults with pertussis are hospitalized or have complications, which could include pneumonia or death.

These diseases are caused by bacteria. Diphtheria and pertussis are spread from person to person through secretions from coughing or sneezing. Tetanus enters the body through cuts, scratches, or wounds.

Before vaccines, as many as 200,000 cases of diphtheria, 200,000 cases of pertussis, and hundreds of cases of tetanus, were reported in the United States each year. Since vaccination began, reports of cases for tetanus and diphtheria have dropped by about 99% and for pertussis by about 80%.

2 Tdap vaccine

Tdap vaccine can protect adolescents and adults from tetanus, diphtheria, and pertussis. One dose of Tdap is routinely given at age 11 or 12. People who did not get Tdap at that age should get it as soon as possible.

Tdap is especially important for healthcare professionals and anyone having close contact with a baby younger than 12 months.

Pregnant women should get a dose of Tdap during every pregnancy, to protect the newborn from pertussis. Infants are most at risk for severe, life-threatening complications from pertussis.

Another vaccine, called Td, protects against tetanus and diphtheria, but not pertussis. A Td booster should be given every 10 years. Tdap may be given as one of these boosters if you have never gotten Tdap before. Tdap may also be given after a severe cut or burn to prevent tetanus infection.

Your doctor or the person giving you the vaccine can give you more information.

Tdap may safely be given at the same time as other vaccines.

3 Some people should not get this vaccine

- A person who has ever had a life-threatening allergic reaction after a previous dose of any diphtheria, tetanus or pertussis containing vaccine, OR has a severe allergy to any part of this vaccine, should not get Tdap vaccine. Tell the person giving the vaccine about any severe allergies.
- Anyone who had come or long repeated seizures within 7 days after a childhood dose of DTP or DTap, or a previous dose of Tdap, should not get Tdap, unless a cause other than the vaccine was found. They can still get Td.
- Talk to your doctor if you:
 - have seizures or another nervous system problem,
 - had severe pain or swelling after any vaccine containing diphtheria, tetanus or pertussis,
 - ever had a condition called Guillain-Barre Syndrome (GBS),
 - aren't feeling well on the day the shot is scheduled.

U.S. Department of Health and Human Services