

**McLaren Print System Order**

**Order No: 48410 Reprint Previous Order No: 47962**  
**Order Date: 2019-09-04**  
**User: Kristin Fudge**  
**Phone: 517-574-9123**

**Ship Location: South Side Medical Center**  
**5525 S. MLKing Jr BLVD**  
**Lansing, Mi 48911**

**Forms**

**Quantity: 100**  
**Paragon Dept No: 68325**  
**Dept Name: South Side Medical Center**  
**Company Number: 810**

**Order Total Price: 0.00**

**Item Number: DCH-0457-A**  
**Item Description: Influenza Vaccine (Live Intranasal) VIS**  
**Revision Date: 8/2019**  
**Print: 2 sided black and white**  
**Paper: 20# White Text**  
**Size: 8.5 x 11**  
**Fold:**  
**Finish: None**  
**Drill: None**  
**Misc Info: Must be ordered with MM-474 Consent**

VACCINE INFORMATION STATEMENT

---

**Influenza (Flu) Vaccine (Live, Intranasal):**  
**What You Need to Know**

Many Vaccine Information Statements are available in Spanish and other languages. For more information, go to [www.cdc.gov/vaccines/imz/](http://www.cdc.gov/vaccines/imz/).

Have an influenza vaccine anytime under supervision or supply to vaccine sites. [www.cdc.gov/vaccines/imz/](http://www.cdc.gov/vaccines/imz/)

**1 Why get vaccinated?**

Influenza vaccine can prevent influenza (flu). Flu is a contagious disease that spreads around the United States every year, usually between October and May. Anyone can get the flu, but it is more dangerous for some people. Infants and young children, people 65 years of age and older, pregnant women, and people with certain health conditions or a weakened immune system are at greater risk of flu complications.

Pneumonia, bronchitis, sinus infections and ear infections are examples of the related complications. If you have a medical condition, such as heart disease, cancer or diabetes, flu can make it worse.

Flu can cause fever and chills, sore throat, muscle aches, fatigue, cough, headache, and runny or stuffy nose. Some people may have vomiting and diarrhea, though this is more common in children than adults.

Each year thousands of people in the United States die from flu, and many more are hospitalized. The vaccine prevents millions of illnesses and flu-related visits to the doctor each year.

**2 Live, attenuated influenza vaccine**

CDC recommends everyone 6 months of age and older get vaccinated every flu season. Children 6 months through 8 years of age may need 2 doses during a single flu season. Everyone else needs only 1 dose each flu season.

Live, attenuated influenza vaccine (called LAIV) is a nasal spray vaccine that may be given to non-pregnant people 2 through 49 years of age. It takes about 2 weeks for protection to develop after vaccination.

There are many flu viruses, and they are always changing. Each year a new flu vaccine is made to protect against those or four viruses that are likely to cause disease in the upcoming flu season. Even when the vaccine doesn't exactly match these viruses, it may still provide some protection.

**3 Talk with your health care provider**

Tell your vaccine provider if the person getting the vaccine:

- Is younger than 2 years or older than 49 years of age.
- Is pregnant.
- Has had an allergic reaction after a previous dose of influenza vaccine, or has any severe, life-threatening allergies.
- Is a child or adolescent 2 through 17 years of age who is receiving aspirin or aspirin-containing products.
- Has a weakened immune system.
- Is a child 2 through 4 years old who has asthma or a history of wheezing in the past 12 months.
- Has taken influenza antiviral medication in the previous 48 hours.
- Cares for severely immunocompromised persons who require a protected environment.
- Is 5 years or older and has asthma.
- Has other underlying medical conditions that can put people at higher risk of serious flu complications (such as lung disease, heart disease, kidney disease, kidney or liver disorders, neurologic or neuromuscular or metabolic disorders).
- Has had Guillain-Barre syndrome within 6 weeks after a previous dose of influenza vaccine.

U.S. Department of Health and Human Services  
Centers for Disease Control and Prevention

800001-01-01