

McLaren Flint  
FLINT, MICHIGAN  
PHYSICIANS ORDERS AND INSTRUCTIONS TO NURSE

**SURGICAL ON – Q  
PAIN PUMP ORDERS**

**1. Fill pump (To be filled by scrub personnel in the OR or LDRP only):**

• MEDICATION:

250 mL 0.5% Bupivacaine (MARCAINE) preservative free injection with 50 mL 0.9% Sodium Chloride preservative free injection

**OR**

250 mL 2% Lidocaine (XYLOCAINE) preservative free injection with 50 mL 0.9% Sodium Chloride preservative free injection

• Flow Rate is not adjustable

• CATHETER:       Single  
                                  Dual

2. Assess patient's pain per protocol

**3. DO NOT REFILL THE PUMP RESERVOIR**

4. Label the pump and attach a tamper proof seal to the fill port

5. Assess site every 4 hours to ensure:

- Tubing in not clamped or kinked
- Site is without redness or drainage
- Dressing is dry and intact
- Flow restrictor is taped to the patient's skin – not in contact with cold or heat therapy
- Air filter is not taped or covered

6. Notify surgeon for / if:

- Patient is experiencing inadequate pain relief
- Excessive drainage noted at the incision or catheter site
- Redness, swelling, tenderness or discharge at the catheter site
- Leakage of local anesthetic from the system
- Patient experiences Hypotension, Palpitations, Bradycardia, Seizure activity, Restlessness, Anxiety, Itching, Nausea or Vomiting
- ON – Q device is empty

7. Discharge Instructions:

- Check with surgeon for removal instructions
- If patient is discharged with the ON – Q pump in place, provide the patient with the ON – Q Patient Guidelines

\_\_\_\_\_  
Physician Signature

\_\_\_\_\_  
Date / Time



PT.

MR#/RM.

DR.