

INTERSTITIAL CYSTITIS FOLLOW UP

PATIENT NAME: _____ Dr. _____ DATE: _____

DOB: _____

URGENCY TODAY: 0 1 2 3 4 5

PAIN TODAY: 0 1 2 3 4 5

MEDICATIONS TAKEN: _____

MEDICATION SIDE EFFECTS: _____

RESULTS WITH MEDICATION: _____

PLAN: _____

RESCUE TREATMENT GIVEN - YES OR NO

Bladder Cocktail:

_____ 30 cc STERILE WATER

_____ 2% LIDOCAINE 10 cc, LOT: _____; EXPIRATION: _____

_____ 8.4% SODIUM BICARBONATE 5 cc LOT: _____; EXPIRATION: _____

_____ 300 mg ELMIRON-SUPPLIED BY THE PATIENT.

_____ Fr CATHETER INSERTED. BLADDER WAS DRAINED. BLADDER COCKTAIL INSTILLED VIA SLOW PUSH INTO THE PATIENT'S BLADDER. CATHETER WAS REMOVED, PATIENT WAS INSTRUCTED TO HOLD THE TREATMENT FOR _____

PATIENT TOLERATED WELL.

Provider Signature: _____ Date: _____ Time: _____

_____ Patient verbalizes understanding of instructions given

Patient Name:

Date of Birth: