STROKE CORE MEASURES	I
DVT Prophylaxis by Day 2	
Antithrombotic by Day 2	
Anticoagulation for history, paroxysmal a fibrillation/flutter	
Thrombolytic Therapy if LKW is within 3-4.5hrs	
Lipid Panel within 48 hours of admission	
Dysphagia Screen (Yale Swallow Screen)	
Intense Statin at discharge (Lipitor 40 mg and up)	*
Antithrombotic at discharge	
Stroke Education (Individualized/address Risk Factors and Prevention/Medication adjustment)	
Rehab (PT/OT/Speech)	
Smoking Cessation	

NIHSS

*If not done, contraindication must be documented by MD, NP or PA.

STROKE

NEURO POWER PLANS

- ED Activated Stroke Alert EKM- use for Stroke Activations in ED
- ED Neuro Deficit- use for non-activated Strokes in ED
- Neuro Inpatient Activated Stroke Alert EKM- use for Rapid Response Code Stroke Activations
- Neuro Ischemic Stroke/TIA with or without alteplase admit or focus orders
- Neuro Ischemic Stroke/TIA admit alteplase (tpa) infusion and follow-up orders
- Neuro Hemorrhagic Stroke admit or focus orders

Admit Orders for Attending; Neurology/Neurosurgery can use Focus Orders

-Make sure we are choosing the appropriate sub phase for each plan when putting in orders.

B-137 (10.22)

BEFAST