

McLaren Print System Order

Order No: 77975
 Order Date: 2023-06-20
 User: Donna Wilson
 Phone: 9898943046

Ship Location: Randy Bowman
 1900 Columbus Ave
 Bay City, MI 48708

Brochures
 Quantity: 1000
 Paragon Dept No: 20200
 Dept Name: PACU
 Company Number: 210

Order Total Price: 192.50

Item Number: BAY-146
 Item Description: Anesthesia Surgery Effects Flyer
 Revision Date: 06/2023
 Print:
 Paper:
 Size:
 Fold:
 Finish:
 Drill:
 Poster:
 Misc Info: SS, Bleed, 80# Offset Text



Spec Info:

SYMPTOMS	WHAT TO DO IF YOU EXPERIENCE THESE SYMPTOMS
NAUSEA / VOMITING	Nausea and vomiting are common for patients recovering from surgery. • Ice chips, saline crackers, or peppermints may help reduce these symptoms. Extreme symptoms may require additional medications, contact your surgeon if not resolved after 24 hours.
TIREDDNESS / FATIGUE	Tiredness and fatigue are common after surgery. This can be due to anesthesia and/or the surgical procedure itself. • Allow yourself extra time to rest and follow activity restrictions as stated by the surgeon/discharge instructions.
SCRATCHY THROAT / HOARSENESS	If a breathing tube is inserted into your mouth, you may experience a scratchy throat and/or hoarseness. • Throat lozenges or gargling with salt water may help. These symptoms generally resolve within 48 hours.
DIZZINESS	Dizziness or mild lightheadedness can be due to medications and/or fluid changes in your system. • Maintain hydration, avoid sudden position changes, and sit for a period at bedside prior to standing to avoid falls.
PAIN	Some ways to control pain include: • Ice packs, splinting incisional area, changing position, taking prescribed pain medications, or over the counter pain relievers. If you have unresolved pain, refer to your discharge instructions and contact your surgeon.

Some patients may notice these symptoms immediately post-operatively, and others may notice symptoms after they are home recovering. If you do experience any of these symptoms, please refer to your discharge instructions for guidance. If symptoms do not improve after 48 hours, please contact your doctor.