

McLaren Print System Order

Order No: 79537  
Order Date: 2023-09-20  
User: Lisa DeWaele  
Phone: 989 894 3906

Ship Location: McLaren Bay Behavioral Health-Attn: Lisa DeWaele  
1900 Columbus Ave.  
Bay City, Michigan 48708

Forms

Quantity: 1000  
Paragon Dept No: 20610  
Dept Name: Behavioral Health  
Company Number: 210

Order Total Price: 33.50

Item Number: BAY-150  
Item Description: Spravato Program Treatment Checklist  
Revision Date: 07/2023  
Print: 1 sided black and white  
Paper: 20# White Text  
Size: 8.5 x 11  
Fold:  
Finish: None  
Drill: 5 Hole Top  
Poster:  
Misc Info: 8.5x11 Black



Spravato Program Treatment Checklist

Please initial upon completion:

- Complete PHQ-9 and GAD-7, report any increase in thoughts.
- Notify provider if patient ate food within the last 2 hours or ingested fluids within the last 30 minutes prior to first dose of medication.
- Vital signs prior to administration, 40 min after 1st dose, and 2 hours after 1st dose.
- Hold medication and notify provider if BP is over 140/90 or below 100/60.
- Monitor sedation level prior to administration, 40 min after 1st dose, and 2 hours after 1st dose.
- Monitor for dissociation: disconnected from self, thoughts, and things around patient prior to administration, 40 min after 1st dose, and 2 hours after 1st dose.
- Observe patient for 2 hours after each treatment.

Provide education:

- Patients should not drive, use heavy machinery, or make financial decisions, not perform any activities that require full alertness until the next day following a restful sleep.
- Provide brief review of common side effects that patients may experience after medication administration. If common side effects occur, they usually happen right after taking Spravato and go away the same day. If prolonged or severe, the patient should discontinue use.
- Review how to properly administer medication with patient.

Discharge criteria:

- Resolution of sedation and dissociation if applicable.
- Patients able to ambulate with and/or as prior to medication administration.
- Vital signs back to patient baseline.
- No signs and symptoms of suicidal behavior.
- Patients must make arrangements to safely get home.

Spec Info:

Signature: \_\_\_\_\_ Initials: \_\_\_\_\_ Title: \_\_\_\_\_

Date: \_\_\_\_\_ Time: \_\_\_\_\_

Signature: \_\_\_\_\_ Initials: \_\_\_\_\_ Title: \_\_\_\_\_

Date: \_\_\_\_\_ Time: \_\_\_\_\_

