McLaren Flint TOTAL JOINT ARTHROPLASTY PRE OP ORDERS

ADMIT TO: Orthopedic Services Surgeon: DIAGNOSIS: Left Right Bilateral	
ALLERGIES:	
HEIGHT: WEIGHT:	
1. SOBTAIN INFORMED CONSENT FOR TOTAL JOINT ARTHROPLASTY	
2. ACTIVITY: As tolerated	
3. ⊠ DIET: NPO after midnight before the day of surgery	
4. ⊠ Surgical prep of operative area morning of surgery to be done in Pre Op Holding area. No perineal shave	
/TE PROPHYLAXIS:	
Apply and Turn ON Sequential Pneumatic Compression (SCD) Device	
Pre-Op Holding (Prior to Induction of Anesthesia)	
	Prophylactic Antibiotics: Administer within 1 hour of surgery start time
Nasal swab for MRSA screen, if not previously done	Cefazolin (Kefzol)
Providine- lodine 5%- apply to each nostril	1 gm if less than 50 kg
If allergic to lodine, Bactroban 2% cream to each nostril if	2 gm if between 50 kg and 120 kg
nasal swab positive for MRSA	☐ 3 gm if more than 120 kg
CMP (Comprehensive Metabolic Profile)	If life-threatening angioedema or anaphylaxis to cephalosporin or
☐ HBA1C (if not obtained within the last six months)	penicillin:
CBC (Complete Blood Count)	Clindamycin 600 mg IVPB – Administer within 1 hour of surgery start time
∑Type and Screen	OR Vancomycin (Administer within 2 hours of surgery start time to run over
Type and Cross 2 units of blood	at least 120 minutes)
MEDICATIONS:	1 gm if less than 100 kg
Dexamethasone (Decadron) 8 mg IV (including patients	1500 mg if 100 kg or more
with Type 1 Diabetes)	If you are using Vancomycin, select all that apply
Pregabalin (LYRICA) 75 mg PO x 1	Cephalosporin allergy or anaphylaxis to penicillin;
☐ Meloxicam (MOBIC) 15 mg PO x 1	MRSA colonization or infection;
	☐ Increase MRSA rate Facility-wide
	Intra-Operative
Irrigation:	Pericapsular Injection:
Intra Operative irrigation of 100,000 units Bacitracin in	Ketorolac 15 mg/ml; Total of 15 mg/volume 1 ml
1000 mL 0.9% NaCl	Morphine 4 mg/ml; Total of 4 mg/volume 1 ml
Intra Operative irrigation of 100,000 units Bacitracin in	Epinephrine 1 mg/ml; Total of 0.6 mg/volume of 0.6 ml
3000 mL 0.9% NaCl	☐ Clonidine 0.1 mg/ml; Total 0.08 mg/0.8 ml☐ Preservative free 0.9% NaClTotal volume of injection of 145 ml
IntraOperative:	Freservative free 0.9 % NaCr Total volume of injection of 145 fill
minutes prior to the end of the procedure	
Tranexamic Acid IV 1000 mg in 0.9% NaCl 100ml x 2. 1	
dose to be given intraoperatively, infuse over 10 minutes.	
1 dose to be given in PACU, 2 hours after first	
dose, infuse at same rate.	
Tranexamic Acid IV 1000 mg in 0.9% NaCl 100ml x 1. 1	
dose to be given topically during the procedure	
(reason:)	

Physician Signature

Date (required)

Time (required)

Page 1 of 1 Revised 4/10/2017 M-1708-224



P

MR.#/P.M.

DR.