


McLaren Print System Order

Order No: 86301 Order Date: 2024-06-17 **Order Request Date: User: Christina Woodman** Phone: 8103421541

Ship Location: McLaren Macomb-North Grove Women's Health Attn: Kim Wisniewski, 44200 Garfield, Suite 164 Clinton Township, Michigan 48038 **Brochures** Quantity: 3 Paragon Dept No: 52043 **Dept Name: Marketing Company Number:**

Order Total Price: 19.50

Item Number: MAC-28 Item Description: WEIGHT LOSS MEDICATIONS AND SURGERY Flyer Pad Revision Date: 06/2024 Print: Paper: Size: Fold: Finish: Drill: Poster: Misc Info: 50 Shts/Pad, 8.5x11 Color SS Bleed 80# Offset Text



Thank you for choosing McLaren Macomb for your surgical needs. We look forward to taking excellent care of you.

To ensure your safety we ask that you stop taking certain medications prior to surgery. Not following these recommendations may lead to your surgery being canceled.

Weight loss medications can increase your risk of neusea and vomiting, increasing your risk of complications during anesthesia. Please look at the list of medications and stop as directed prior to your surgery.

- Weight loss medications that must be stopped seven days before surgery:
 • Exervatide (Bydureon)
 • Transpatide/Zepbound (Morjaro)

 • Semagluide (Ozempic, Wegovy, and Rybelau)
 • Transpatide/Zepbound (Morjaro)

- Linglutide (Savenda, Victora)
 Dulaglutide (Trulicity) Weight loss medications that must be stopped the day of surgery:
- · Orlatat (Kanical, A5)
- PEG-Lovenatide
 Phontermine (Adjuer P) Empegifiscin Gardianon)
- Weight loss medications that must be stopped 24 hours before surgery: Exercicle (Byetta)
- If you have any questions, please call your surgeon for clarification

Spec Info:



For questions please call your surgeon mclanen.org