

**McLaren Print System Order**

Order No: 89344  
Order Date: 2024-10-29  
Order Request Date:  
User: Nick VanHorn  
Phone: 989-894-6458

Ship Location: McLaren Bay Region Attn: Nick VanHorn  
300 Mulholland St  
Bay City, 48708

Brochures  
Quantity: 500  
Paragon Dept No: 20200  
Dept Name: Marketing  
Company Number:

Order Total Price: 96.25

Item Number: BAY-113  
Item Description: DIABETES, WEIGHT LOSS MEDICATIONS, AND SURGERY Flyer  
Revision Date: 10/2024  
Print:  
Paper:  
Size:  
Fold:  
Finish:  
Drill:  
Poster:  
Misc Info: 8.5x11 SS Bleed 80# Offset



Thank you for choosing McLaren Bay Region for your surgical needs. We look forward to taking excellent care of you.

To ensure your safety we ask that you stop taking certain medications prior to surgery. Not following these recommendations may lead to your surgery being cancelled.

Weight loss medications can increase your risk of nausea and vomiting, increasing your risk of complications during anesthesia. Please look at the list of medications and stop as directed prior to your surgery.

Weight loss medications that must be stopped seven days before surgery:

- Exenatide (Bydureon)
- Semaglutide (Ozempic, Wegovy, and Rybelsus)
- Liraglutide (Saxenda, Victoza)
- Dulaglutide (Trulicity)

- Tirzepatide/Zepbound (Mounjaro)
- Lixisenatide (Aduvia)
- Abiraterone (Zytiga)
- PEG-Losartan
- Phentermine (Adipex-P)

Weight loss medications that must be stopped 72 hours prior to surgery:

- Orlistat (Xenical, Alli)
- Empagliflozin (Jardiance)
- Exenatide (Byetta)

If you have any questions, please call your surgeon for clarification.

**Spec Info:**